

Détail des stations

- Le Moura, forêt communale de Tournay

- Tournay (65)

Le 20 juillet 2014, Gilles Corriol ; Robert Cazenave ; Germaine Dubranat ; AMB

Source Prospection

<u>GC1</u>	54 taxons	Amanita rubescens, Amanita rubescens f. annulosulfurea, Boletus aereus, Boletus aestivalis, Cantharellus friesii, Cantharellus subpruinosis, Chlorociboria aeruginascens, Climacodon pulcherrimus, Collybia dryophila, Collybia fusipes, Collybia inodora (Micromphale inodorum (Pat.) Svrcek), Collybia peronata, Cortinarius bolaris, Creopus gelatinosus, Daedalea quercina, Daedaleopsis confragosa, Diatrype decorticata, Gyroporus castaneus, Hymenochaete rubiginosa, Hypoxylon multifforme, Laccaria tortilis, Lactarius piperatus, Leccinum carpini, Leccinum crocipodium, Leccinum quercinum, Marasmiellus candidus, Marasmiellus ramealis, Marasmius rotula, Megacollybia platyphylla, Mycena pura, Mycoacia uda (Phlebia uda (Fr.) Nakasone), Oudemansiella radicata, Phlebiella vaga (Phlebiella sulphurea (Pers. : Fr.) Ginns & Lefebvre), Polyporus lentus (Polyporus tuberaster (Jacq.) Fr.), Polyporus varius var. nummularius (Polyporus leptocephalus var. nummularius (Bull. : Fr.) Courtec.), Psathyrella laevisissima, Rickenella fibula, Russula amoena, Russula aurora, Russula cyanoxantha, Russula puellula, Russula risigallina, Russula virescens, Schizopora paradoxa, Scleroderma citrinum, Sebacina incrustans, Simocybe centunculus (Ramicola centunculus (Fr. : Fr.) Watl.), Simocybe sumptuosa (Ramicola sumptuosa (Orton) Watl.), Skeletocutis nivea, Steccherinum ochraceum, Stereum hirsutum, Strobilomyces strobilaceus (Strobilomyces floccopus (Vahl. : Fr.) P. Karst.), Trametes versicolor, Xerocomus rubellus.
<u>Altitude</u>	280 - 290	
<u>Topographie</u>		
<u>Type de relief</u>		
<u>Pente</u>		
<u>Exposition</u>		
<u>Roche mère</u>		

Groupement végétal